



PI-C3017 V1.0

Product Name

Name: RPMI 1640 Medium, without L-Glutamine

Cat. No.: C3017-0500

Size: 500 mL

Product Description

RPMI 1640 Medium without L-Glutamine has been specifically developed for the long-term culture of blood cells, the culture of normal and abnormal human leukocytes (e.g., neoplastic WBC's) and is now used as a general medium (with serum) for hybridoma cultures. Roswell Park Memorial Institute (RPMI) 1640, when properly supplemented, has demonstrated wide applicability for supporting the growth of many types of cells in culture, including human lymphocytes.

Most common types of media consist of an isotonic, buffered basal nutrient-enriched environment which provides an energy source, inorganic salts, vitamins, amino acids as well as additional constituents (e.g., supplements) according to the demands of a particular cell line. This relatively more complex medium formulation provides the optimal cell culture environment which mimics those of the in vivo environment including basic nutritional requirements, osmotic pressure, physiological pH, and temperature among other considerations. At a minimum, it consists of the foundation medium components that are all part and parcel of a pre-tested complete media to assist the cells in meeting their metabolic demands.

RPMI 1640 medium contains no growth-promoting factors or antimicrobials.

L-Glutamine, a precursor of glutamate, is one of the most readily available sources of energy for many rapidly dividing cell types for use in vitro and is a key component and essential amino acid that is required in many cell culture media formulations and in virtually all mammalian cells in culture. Sodium pyruvate serves as an additional and easily accessible carbohydrate energy source for cells in culture. Along with D-glucose, these balanced energy sources serve as carbon skeletons for cell anabolic processes in addition to nucleic acid metabolism and protein production while limiting the potential accumulation of the toxic ammonia gas.

Composition

| Ingredients | mg/L | Ingredients | mg/L |
|---------------------------------------|---------|------------------------------------|----------|
| INORGANIC SALTS | | | |
| Calcium nitrate tetrahydrate | 100.000 | Sodium chloride | 6000.000 |
| Magnesium sulphate anhydrous | 48.840 | Sodium phosphate dibasic anhydrous | 800.000 |
| Potassium chloride | 400.000 | | |
| AMINO ACIDS | | | |
| Glycine | 10.000 | L-Leucine | 50.000 |
| L-Arginine hydrochloride | 241.000 | L-Lysine hydrochloride | 40.000 |
| L-Asparagine | 50.000 | L-Methionine | 15.000 |
| L-Aspartic acid | 20.000 | L-Phenylalanine | 15.000 |
| L-Cystine dihydrochloride | 65.200 | L-Proline | 20.000 |
| L-Glutamic acid | 20.000 | L-Serine | 30.000 |
| L-Histidine hydrochloride monohydrate | 20.960 | L-Threonine | 20.000 |
| L-Tryptophan | 5.000 | L-Tyrosine disodium salt | 28.830 |



RPMI 1640 Medium product information

| Y | | | PI-C3017 V1.0 |
|--------------------------|----------|-----------------------------|---------------|
| L-Hydroxyproline | 20.000 | L-Valine | 20.000 |
| L-Isoleucine | 50.000 | | |
| Vitamins | | | |
| Choline chloride | 3.000 | Riboflavin | 0.200 |
| D-Biotin | 0.200 | Thiamine hydrochloride | 1.000 |
| D-Ca-Pantothenate | 0.250 | Vitamin B12 | 0.005 |
| Folic acid | 1.000 | i-Inositol | 35.000 |
| Niacinamide | 1.000 | p-Amino benzoic acid (PABA) | 1.000 |
| Pyridoxine hydrochloride | 1.000 | | |
| OTHERS | | | |
| D-Glucose | 2000.000 | Phenol red sodium salt | 5.300 |
| Glutathione reduced | 1.000 | Sodium bicarbonate | 2000.000 |

Storage and Stability

The product should be kept at 2 - 8°C.

The product is **light-sensitive** and therefore should not be left in the light.

Shelf life: 12 months from date of manufacture.

Procedure

- 1. Take a bottle from the refrigerator at 2-8°C and read the label.
- 2. Wipe the outside of the bottle with a disinfectant solution such as 70% ethanol.
- 3. Pipette appropriate volume using aseptic/sterile technique under a laminar-flow culture hood.
- 4. Add 2 mM of L-Glutamine 200 mM Solution. L-Glutamine, a precursor of glutamate, is one of the most readily available sources of energy for many rapidly dividing cell-types for use in vitro and is a key component and essential amino acid that is required in many cell culture media formulations and in virtually all mammalian cells in culture.
- 5. Add antibiotics or other nutrients if desired.

Quality control

RPMI 1640 Medium, without L-Glutamine is tested for sterility, pH, osmolality, and endotoxin concentration. In addition, each batch is tested for cell growth performance.

Manufacturer

Shanghai Dr. Cell Co., Ltd.

Issue Date

June 2023









PI-C3017 V1.0

Precaution and Disclaimer

For research use only, not for clinical diagnosis, and treatment.

